

FIRST-radio sources in clusters of galaxies

A.G. Gubanov, V.P. Reshetnikov

Astronomical Institute of St.Petersburg State University, Russia

Abstract

The detailed identifications of the FIRST and NVSS radio sources with optical objects in APM and in DSS surveys are carried out for 26 rich Abell clusters of galaxies. 99 radio sources are identified with optical objects, 40 have probable identifications, and 187 are not identified from 326 radio sources in the considered fields (within Abell radius of each cluster of galaxies). 20 radio sources are definitely in clusters and 34 can be clusters members with high confidence. Therefore, $\approx 30\text{-}40\%$ of the FIRST radio sources can be identified with optical objects on the base of APM and of DSS data. On average, one can find 2 identified sources per cluster.

In the framework of previously announced project (A.G.Gubanov et al. 1997, in “Problems of modern radioastronomy”, XXVII Radio Astronomy conference, St.Petersburg, V.1, P.168), we present detailed identifications of the FIRST (and NVSS) radio sources with objects from APM and DSS surveys for 26 rich Abell clusters of galaxies (Table 1).

Identifications are performed for all radio sources within Abell radius of cluster of galaxies (3 Mpc for the Hubble constant of 50 km/s/Mpc). The clusters selected within the processed zone of the FIRST survey are part of a larger sample of rich clusters (with richness ≥ 2) and of clusters containing cD-galaxies.

Table 1. List of clusters.

ACO	RA(J2000)			DEC(J2000)			z	ACO	RA(J2000)			DEC(J2000)			z
	h	m	s	o	'	"			h	m	s	o	'	"	
A0586	07	32	17.8	+31	37	34	0.1710	A1033	10	31	33.8	+35	04	34	0.1258
A0642	08	19	05.7	+30	02	34	0.2063e	A1035	10	32	07.3	+40	12	33	0.0785
A0690	08	39	14.3	+28	51	24	0.0788	A1068	10	40	47.2	+39	58	20	0.1375
A0705	08	47	38.7	+30	00	56	0.1138e	A1073	10	42	26.6	+36	38	17	0.1390
A0715	08	54	44.4	+35	24	33	0.1685e	A1081	10	44	49.5	+35	34	14	0.1585
A0727	08	59	13.1	+39	26	19	0.0992e	A1094	10	47	32.8	+27	31	10	0.2004
A0781	09	20	23.2	+30	26	16	0.1763e	A1120	10	53	15.5	+30	48	02	0.2218e
A0800	09	28	30.0	+37	47	53	0.2223e	A1175	11	09	13.7	+33	10	44	0.2487
A0812	09	32	35.4	+37	53	42	0.1452e	A1178	11	09	50.1	+34	35	43	0.2596
A0908	09	59	35.8	+22	25	36	0.2026e	A1182	11	10	19.0	+31	46	43	0.1660e
A0943	10	12	12.2	+33	37	10	0.1485e	A1190	11	11	46.2	+40	50	42	0.0763
A0961	10	16	29.5	+33	37	01	0.1241	A1198	11	12	48.0	+30	22	40	0.1660e
A0963	10	17	09.7	+39	01	00	0.2060	A1258	11	26	08.7	+25	26	30	0.1469e

We obtained the following results:

1. Among 326 radio sources in the considered fields of 26 clusters (and within Abell radius), 99 radio sources are confidently identified with optical objects in the APM and DSS surveys, 40 sources have probable identifications, and 187 are not identified.

Among identified radio sources, 20 are definitely in clusters and 34 can be clusters members with high probability.

Therefore, the expectations to identify $\approx 30\text{-}40\%$ FIRST radio sources and to find, on average, 2 sources per cluster are coming true.

One can expect to obtain ≈ 8000 identifications of the galaxies in ACO-clusters with the FIRST-type radio sources in all Abell clusters.

2. Only 11 radio sources consist of one component (within the FIRST-survey resolution of about $5''$) and only 5 are compact (with diameter $\leq 5''$) among 20 definitely lying in clusters

sources.

Morphology of identified radio sources is typical for clusters radio-galaxies:

5 have WAT structure;

5 – HT;

5 - extended FR I-type radio sources (with uncertain morphological type);

5 - compact.

Centers of radio sources are often essentially displaced from the positions of host galaxies in our sample and, therefore, visual analysis of radio-optical maps is needed for correct identification. Graphical interface of the database of clusters created at the Astronomical Institute of the SPbSU (A.G.Gubanov, V.B.Titov, 1997, in “Problems of modern Radio Astronomy”, XXVII Radio Astronomy conference, St.Petersburg, V.1, P.320) is an excellent and useful tool for such purposes (URL [**http://future.astro.spbu.ru/Clusters.html**](http://future.astro.spbu.ru/Clusters.html)).

3. In most cases the identified radio sources have radio luminosities, $\log(L_r \text{ [erg/sec]})$, typical for radio galaxies – 40.5-42 within frequency diapason 10 MHz – 10 GHz and are identified with bright elliptical galaxies.

4. It seems interesting that dynamically active clusters (consisting of subclusters of comparable richness) often contain several radio galaxies with comparable power (for instance, A1033, A1035, A1068, A1081, A1190, A1258). This is not typical for more regular clusters.

5. It should be noted that the flux densities in the FIRST, NVSS catalogues and apparent magnitudes in the APM-survey are contain often significant systematic and random errors and cannot be useful for correct statistical analysis. Careful individual reduction of the data of digitized surveys and special measurements are highly needed.

Several examples of identified radio sources are presented in Fig.1–3.

This research is supported by the Federal Programme “Integration” (project *N* 578) and by the Russian Foundation for Basic Research (grant *N* 97-02-18212).




Figure 1: One from two brightest galaxies in cluster center, HT radio galaxy. Faint compact detail in the tail of this radio source coincides in position with other cluster galaxy (i.e. possible cluster radio galaxy too).




Figure 2: The pair of close galaxies, probably, eastern is radio galaxy.




Figure 3: HT radio source is identified with second bright galaxy in cluster center.